

Research Article

Independent Determinants of Urinary Albumin Excretion and Confounding Variables in Type 2 Diabetic Patient

Sunita Neupane¹, Sujata Shrestha^{2*}¹Department of Biochemistry, Bhaktapur District Hospital, Nepal²Department of Biochemistry, Paropakar Maternity Hospital, Nepal

Copyright: © 2016 Sujata Shrestha, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Background: Microalbuminuria is a known risk factor for the development of clinical nephropathy in diabetes and also an independent risk factor for cardiovascular disease. Microalbuminuria is a marker of a pathophysiological process that causes both increased renal albumin loss and atherothrombosis. Microalbuminuria is hallmark for early detection of diabetic nephropathy. An elevated urinary albumin excretion is a marker of endothelial dysfunction that symbolizes the kidney's way to translate the existence of vascular damage. The aim of this study was to evaluate the independent determinants of urinary albumin excretion, and association between biochemical parameters and socio-demographic factors in Diabetic patients.

Materials and Methods: This is a hospital based cross sectional study included diagnosed case of Diabetic patients. Serum uric acid concentrations were measured by enzymatic method (uricase-peroxidase), HbA1c was measured using the principle of dry chemistry, Blood Sugar measured by Glucose oxidase peroxidase (GOD/POD) method and urinary albumin excretion was measured with an immunoturbidometric assay.

Results: Based on categorization of Urinary albumin excretion, 65% normoalbuminuric, 27% microalbuminuric and 8% macroalbuminuric are found in our study population. The frequency of hyperuricemia was found to be 43%. The prevalence of albuminuria increased significantly with increasing glycaemia. Pearson's Correlation coefficient by bivariate analysis of Urinary albumin excretion with confounding variables shows significant positive correlation with onset of DM ($r=0.203$, $p=0.013$), systolic blood pressure ($r=0.355$, $p=0.001$), diastolic blood pressure ($r=0.405$, $p=0.001$), uric acid ($r=0.352$, $p=0.001$), hba1c ($r=0.212$, $p=0.005$) and smoking ($r=0.265$, $p=0.01$). Multiple regression test shows that independent determinant of UAE are blood pressure {Diastolic ($\beta=0.313$, $P=0.006$) / Systolic ($\beta=0.309$, $P=0.002$)}, hba1c ($\beta=0.187$, $P=0.010$), Uric acid ($\beta=0.331$, $P=0.0001$) and Onset of DM ($\beta=0.199$, $P=0.041$).

Conclusion: Albuminuria is therefore an important risk factor to measure in patients at risk. The findings extend the relationship between confounding variables and the urinary albumin excretion which emphasize on the importance of screening for microalbuminuria, Serum Uric Acid to prevent renal dysfunction, HbA1c measurement on a regular interval for good glycemic control and the other variables for regular physiological process of body. Further examination is needed in a large population size to clarify the validity between the biochemical parameters

Key words: Urinary albumin excretion, Uric acid, Diabetes Mellitus, HbA1c

Introduction

Microalbuminuria was first defined by Mogensen termed as a moderate increase in the level of urine albumin. It occurs when the kidney leaks small amounts of albumin into urine, in other words, when there is an abnormally high permeability for albumin in the renal glomerulus. [1] Article which are published in Pubmed are mostly taken for reference materials. Microalbuminuria, in addition to being the hallmark for early detection of diabetic nephropathy, is a marker for endothelial dysfunction and a known risk factor for cardiovascular events and mortality in patients with Diabetes Mellitus (DM). [2] Nephropathy was graded as follows: normoalbuminuria, urinary albumin excretion less than 30 mg/g Cr; microalbuminuria, 30 to 300 mg/g Cr; or macroalbuminuria, more than 300 mg/g Cr. [3]

Microalbuminuria is an important adverse predictor of glycemic outcomes in pre-diabetes. Prediabetic patient with increased microalbuminuria even in the normal range is associated with increased progression to diabetes and decreased reversal to normoglycemia. So, Prediabetic patient with microalbuminuria needs more study and discovery to prevent diabetes in them. [4] Cardiovascular atherosclerotic disease is seen in non-diabetic as well as in diabetic patients having elevated urinary albumin excretion

(UAE). [5,6] Microalbuminuria is also an integral component of the metabolic syndrome characterized by insulin resistance. [7,8] The aim of this study was to evaluate the independent determinants of urinary albumin excretion, association between biochemical parameters and socio-demographic factors in Diabetic patients.

Materials and Methods

A cross-sectional study was done in Bhaktapur district hospital with collaboration of internal medicine and ward in 150 patients who are diagnosed as diabetic based on the WHO criteria. Patient fulfilling the criteria of research work were included in the study population whereas Patient who are well diagnosed of kidney failure, organ transplant, chronic alcoholic, using diuretics, and other drug which lowers uric acid were excluded. Serum uric acid (SUA) concentrations were measured by enzymatic method (uricase-peroxidase), Glycated Hemoglobin (HbA1c) was measured using the principle of dry chemistry, Blood Sugar measured by Glucose Oxidase/Peroxidase (GOD/POD) method and UAE was measured

***Corresponding author:** Ms Sujata Shrestha, B.Sc. MLT, Department of Biochemistry, Paropakar Maternity Hospital, Nepal, Tel: +977-9841671000; Fax: +977-9841671000; E-mail: ssuju.046@gmail.com

Received: October 10, 2016; **Accepted:** November 21, 2016; **Published:** November 24, 2016

with an immunoturbidometric assay. During the data collection process, after taking informed consent from patient, proforma was fill up that includes patient history, Previous records, biochemical parameters as well as other socio demographic variables like smoking, dietary habit, Current treatment, Blood pressure, alcoholism, Height, Weight etc for further analysis. Random urine sample was collected for urine microalbumin test whereas 12 hr fasting for fasting blood sugar measurement, 2 hr after meal for Post Prandial(PP) sugar. EDTA blood used for the HbA1c test and random blood for SUA test.

Statistical analysis

Data were expressed as the means \pm SD, Correlations were performed with the Pearson's tests, depending on the distribution of variables. Multiple linear regressions were performed with UAE as the dependent variable using the SPSS version 20. *P* values <0.05 (two tailed) were considered to be significant.

Results

Table 1 shows the occurrence of the diabetic patient as well as the socio demographic variables including biochemical parameters. Age group >25 and <80 years were taken for the research purpose in 150 Diabetic patients. Population who were having DM <10 years was found to be 65% whereas duration >10 years was 35%. In our study population, majority was male (72%, $n=108$) and female (28%, $n=42$). Most of the patient was obese with their BMI >25 kg/ m^2 . When we measured the SBP and DBP, about 60% were within the normal range. Past smokers with past alcoholic were about 55% in this study population. Based on the treatment modality, majority of the patient were under OHA (76%) with non-vegetarian diet (95%). The frequency of hyperuricemia was found to be 43%. Based on the categorization of urinary albumin excretion, 65% ($n=98$) were normoalbuminuric, 27% ($n=40$) microalbuminuric and 8% ($n=12$) macroalbuminuric. 20% of the population were under good glycemic control and 80% under the poor glycemic control. About 75% of DM patient were beyond the normal range of blood sugar.

Table 2 presents the biochemical parameters and socio demographic variables in descriptive statistics form i.e. mean and Standard Deviation.

Table 3 shows the Positive correlation of Urinary albumin excretion with all the variables. From the above table, UAE is significantly correlated with onset of DM ($r=0.203$, $P=0.013$), Systolic Blood Pressure ($r=0.355$, $P=0.001$), Diastolic Blood Pressure ($r=0.405$, $P=0.001$), Serum Uric acid ($r=0.352$, $P=0.001$), HbA1c ($r=0.212$, $P=0.005$) and Smoking ($r=0.265$, $P=0.01$).

Table 4 shows independent determinant of UAE in T2DM patient. Multiple regression test shows that the independent determinant of UAE are DBP ($\beta=0.313$, $P=0.006$), SBP ($\beta=0.309$, $P=0.002$), HbA1c ($\beta=0.187$, $P=0.010$), SUA ($\beta=0.331$, $P=0.0001$) and Onset of DM ($\beta=0.199$, $P=0.041$).

Discussion

The cross-sectional study has been carried on 150 T2DM patients. Study includes all age group from age 25 to 80 years old which shows that the occurrence of kidney dysfunction does not limit for specific age group, onset, and duration of DM. The lack of correct treatment on time and nonspecific test may disturb the physiological process of body. The American Diabetes Association recommends that patients with type 2 diabetes be tested for albuminuria at the time of initial diabetes diagnosis and yearly thereafter. [9] The prevalence of albuminuria increased significantly with increasing glycaemia. Our study shows that UAE is positively correlated with onset of DM, Blood pressure, SUA, HbA1c, smoking, age, BMI, Fasting sugar, PP sugar and other variables. Likewise, several studies

show that albuminuria was also significantly correlated with indices of glycaemic control, i.e. HbA_{1c}, fasting glucose and duration of diabetes, as well as insulin treatment as a categorical variable. Renal lesions are seen in the patients who are under insulin treatment with longer duration of diabetes. [10] BMI and the categorical variables of elevated waist circumference and the metabolic syndrome according to NCEP-ATP-III criteria were correlated with albuminuria, which also supports our study from several regards. [11] This observation is again in line with previous communications and the observation that UAE reflects insulin resistance. [12] Tseng CH found a positive correlation between SUA and albuminuria in male T2DM in whom confounding factors had been excluded, such as use of allopurinol, diuretics or alcohol consumption. [13] Also, Neupane S et al found the positive association between SUA, microalbumin and glycated hemoglobin in T2DM patients. [3] One study has shown that metabolic syndrome is associated with both microalbuminuria and chronic kidney disease [14] and the same is observed in hypertensive patients. [15] Increased UAE brings diabetic nephropathy, and high blood pressure to glomerulopathy, especially if diabetic retinopathy exists [16]. The relative contribution of SBP and DBP varied from one country group to another, the contribution was proportional to the mean SBP/DBP values observed within each country group. Plasma glucose was also an important and expected contributor to high urinary albumin. In those country groups where plasma glucose was an independent contributor, so was diabetic retinopathy. This finding is in close accordance with the data in the UKPDS [17]. In this study, we were unable to include insulin levels and lipid profile [18, 19] that may act as preliminary indicator for cardiovascular risk induced due to high UAE in T2DM patients. Alcohol intake is also correlated with high UAE which contrasts with a Renaud S et al report. [20] Quality and quantity of alcohol intake perhaps protect from premature mortality [21, 22] but over alcohol intake lead high blood pressure. [23] Considering these variables, alcohol intake, smoking, dietary etc did not contribute independently to high urinary albumin in our study population. An elevated UAE is a marker of endothelial dysfunction that symbolizes the kidney's way to translate the existence of vascular damage. Renal disease is both a cause and consequence of hypertension. Reduction of blood pressure and albuminuria reduces cardiovascular risk and renal risk. Although more frequent in diastolic than systolic dysfunction, it appears to indicate a worse prognosis in the latter class. Albuminuria is frequently associated with diabetes, and indicates a worse prognosis in patients with heart failure and diabetes than in those with heart failure alone. This is probably due to the bidirectional relationship between diabetes and heart failure.

Conclusion

Albuminuria is therefore an important risk factor to measure in patients at risk. The findings extend the relationship between confounding variables and UAE excretion which emphasize on the importance of screening for microalbuminuria and SUA to prevent renal dysfunction, HbA1c measurement on a regular interval for good glycemic control and the other variables for regular physiological process of body. Further examination is needed in a large population size to clarify the validity between the biochemical parameters

Conflicts of Interest

None Declared

Acknowledgement

I would like to appreciate Mr Prasadha Raj Neupane and Mr Ram Prasad Acharya for their technical support during this research. Also, want to thank all of my teachers for their constant encouragement and motivation. We hereby also declare that there is no any financial support or relationships that may pose conflict of interest.

Table 1: Frequency of Variables.

Variables	Frequency(n=150)
Duration of DM(<10 years) (>10 years)	65% 35%
Sex(Male/Female)	(72% / 28%)
Body Mass Index(BMI) (<20 / >20) kg/m ²	25% / 75%
Systolic Blood Pressure(SBP) (100-140)mmHg (>140 mmHg)	60% 40%
Diastolic Blood Pressure(DBP) (60-90)mm Hg (>90 mm Hg)	65% 35%
Smoking(Non/ Past/ Current)	(40% / 54% / 6%)
Alcoholism(Non/ Past/ Current)	(40% / 56% / 4%)
Current treatment (Diet / OHA/ Insulin)	(2% / 76% / 22%)
Diet(vegetarian/Non-veg)	(5% / 95%)
Serum Uric Acid(<6.5 mg/dl) (>6.5 mg/dl)	57% 43%
Urinary albumin Excretion (normoalbuminuric) (microalbuminuric) (macroalbuminuric)	65% 27% 8%
HbA1c (<6.4%) (>6.4%)	20% 80%
Fasting Blood Sugar (60-110)mg/dl (>110 mg/dl)	26% 74%
Post Prandial Blood Sugar (100-140)mg/dl (>140 mg/dl)	23% 77%

*OHA: Oral Hypoglycemic Agent

Table 3: Pearsons Correlation Coefficient by Bivariate Analysis of Urinary Albumin Excretion with Confounding Variables

Variables	r	p
Onset of DM	0.203	0.013*
Systolic Blood Pressure	0.355	0.001*
Diastolic Blood Pressure	0.405	0.001*
Serum Uric acid	0.352	0.001*
HbA1c	0.212	0.005*
Smoking	0.265	0.001*
Age	0.119	0.146
BMI	0.028	0.391
Alcoholism	0.033	0.693
Diet	0.117	0.153
Fasting Blood Sugar	0.024	0.774
Post prandial Blood Sugar	0.007	0.930

*. Correlation is significant at the 0.05 level (2-tailed).

Table 2: Patient Characteristics Using Descriptive Statistics

Variable	Mean	Std. Deviation
Age	57.273	12.0713
Duration of DM	7.687	6.6717
Onset of DM	49.213	10.2359
Body Mass Index(BMI)	23.5104	3.04382
Systolic Blood Pressure	140.000	21.6980
Diastolic Blood Pressure	92.133	15.0430
Serum Uric Acid	6.393	1.4061
Urinary albumin Excretion	55.865	75.9930
HbA1c	8.269	2.0572
Fasting Blood Sugar	147.460	49.0195
Post Prandial Blood Sugar	207.493	91.0013

References

- Mogensen CE (1984) Microalbuminuria predicts clinical proteinuria and early mortality in maturity-onset diabetes. *N Engl J Med* 310: 356-360. [crossref]
- Newman DJ, Mattock MB, Dawney AB, Kerry S, McGuire A, et al. (2005)

Table 4: Independent Determinant of Urinary Albumin Excretion in T2DM Patient

variable	β	p
Diastolic Blood Pressure	0.313	0.006
Systolic Blood Pressure	0.309	0.002
HbA1c	0.187	0.010
Serum Uric Acid	0.331	0.0001
Onset of DM	0.199	0.041

Systematic review on urine albumin testing for early detection of diabetic complications. *Health Technol Assess* 9: 160–163.

- Neupane S, Dubey RK, Gautam N, Agrawal KK, Jayan A, et al. (2016) Association between serum uric acid, urinary albumin excretion, and glycated hemoglobin in Type 2 diabetic patient. *Nigerian Medical Journal: Journal of the Nigeria Medical Association* 57: 119-123.
- Dutta D, Choudhuri S, Mondal SA, Mukherjee S, Chowdhury S, et al. (2015) “Urinary albumin?:creatinine ratio predicts prediabetes progression to diabetes and reversal to normoglycemia: role of associated insulin resistance, inflammatory cytokines and low vitamin D”. *Journal of Diabetes* 6: 316–322.
- Deckert T, Yokoyama H, Mathiesen E, Rønn B, Jensen T, et al. (1996) Cohort study of predictive value of urinary albumin excretion for atherosclerotic vascular disease in patients with insulin dependent diabetes. *BMJ* 312: 871–874.
- Dinneen SF, Gerstein HC (1997) The association of microalbuminuria and mortality in non-insulin-dependent diabetes mellitus. A systematic overview of the literature. *Arch Intern Med* 157: 1413-1418. [crossref]
- McFarlane SI, Banerji M, Sowers JR (2001) Insulin resistance and cardiovascular disease. *J Clin Endocrinol Metab* 86: 713-718. [crossref]
- American diabetes association position statement: Diabetic nephropathy. *Diabetes Care* 2003; 26: S94–S98
- American Diabetes Association (2005) Standards of medical care in diabetes. *Diabetes Care* 28 Suppl 1: S4-S36. [crossref]
- Ritz E, Viberti GC, Ruilope LM, Rabelink AJ, Izzo JL Jr, et al. (2010) Determinants of urinary albumin excretion within the normal range in patients with type 2 diabetes: the Randomised Olmesartan and Diabetes Microalbuminuria Prevention (ROADMAP) study. *Diabetologia* 53: 49-57.
- Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (2001) Executive Summary of The Third Report of The National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, And Treatment of High Blood Cholesterol In Adults (Adult Treatment Panel III). *JAMA* 285: 2486-2497. [crossref]
- Parvanova AI, Trevisan R, Iliev IP, Dimitrov BD, Vedovato M, et al. (2006) Insulin resistance and microalbuminuria: a cross-sectional, case-control study of 158 patients with type 2 diabetes and different degrees of urinary albumin excretion. *Diabetes* 55: 1456-1462. [crossref]
- Tseng CH (2005) Correlation of uric acid and urinary albumin excretion rate in patients with type 2 diabetes mellitus in Taiwan. *Kidney Int* 68: 796-801. [crossref]
- Chen J, Muntner P, Hamm LL, Jones DW, Batuman V, et al. (2004) The metabolic syndrome and chronic kidney disease in U.S. adults. *Ann Intern Med* 140: 167-174. [crossref]
- Pascual JM, Rodilla E, Gonzalez C, Pérez-Hoyos S, Redon J (2005) Long-term impact of systolic blood pressure and glycemia on the development of microalbuminuria in essential hypertension. *Hypertension* 45: 1125-1130. [crossref]
- Osterby R, Gall MA, Schmitz A, Nielsen FS, Nyberg G, et al. (1993) Glomerular structure and function in proteinuric type 2 (non-insulin-dependent) diabetic patients. *Diabetologia* 36: 1064–1070.
- Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33) (1998). UK Prospective Diabetes Study (UKPDS) Group. *Lancet* 352: 837-853. [crossref]
- Marre M, Bouhanick B, Berrut G (1994) Microalbuminuria. *Curr Opin Nephrol Hypertens* 3: 558-563. [crossref]
- Metcalf PA, Baker JR, Scott AJ, Wild CJ, Scragg RKR, et al. (1992) Albuminuria in people at least 40 years old: effect of obesity, hypertension, and hyperlipidemia. *Clin Chem* 38: 1802–1808.

20. Renaud S, de Lorgeril M (1992) Wine, alcohol, platelets, and the French paradox for coronary heart disease. *Lancet* 339: 1523-1526. [\[crossref\]](#)
21. Metcalf PA, Baker JR, Scragg RKR, Dryson E, Scott AJ, et al. (1993) Albuminuria in people at least 40 years old: effect of alcohol consumption, regular exercise, and cigarette smoking. *Clin Chem* 39:1793-1797.
22. Grønbaek M, Deis A, Sørensen TI, Becker U, Schnohr P, et al. (1995) Mortality associated with moderate intakes of wine, beer, or spirits. *BMJ* 310: 1165-1169. [\[crossref\]](#)
23. Lang T, Degoulet P, Aime F, Devries C, Jacquinet-Salord MC, et al. (1987) Relationship between alcohol consumption and hypertension prevalence and control in a French population. *J Chronic Dis* 40: 713-720.